ProClova® XL Herbicide

ProClova XL is a new exciting clover-safe herbicide that kills key weeds in newly sown leys and established grassland.

ProClova XL:

- Has a wide application window from March to October
- Kills key weeds in newly sown leys, notably chickweed, fat-hen, redshank and seedling docks
- Delivers the highest and most reliable level of control of docks in established grassland where clover preservation is key
- Can be used on grass leys that contain red clover if applied in the year after establishment. (Do NOT use on year One of Red Clover establishment.)

Benefits

- Use ProClova XL on newly sown leys to ensure grass and clover establish and dominate the sward
- Improve the productivity of established grass and clover swards so that you have more and better quality forage
- Also benefit from great control of other problem weeds buttercups, dandelions, hogweed and cow parsley

Application

- Always apply ProClova with XL adjuvant
- Target weeds at the correct growth stage, if you have topped the weeds wait for sufficient regrowth before spraying
- Apply with at least 75% drift reducing nozzles
- DO NOT apply this product to land parcels where an intersecting watercourse is present
- When applying to land parcels where other types of watercourses are present, respect an unsprayed buffer zone of 3m to these watercourses
- Apply to actively growing weeds, ensure applications are made where temperatures exceed 8°C for three days prior to spraying and 3 days post spraying
- 1 application per calendar year
- Apply with 200 L water / hectare

Bandon Co-Operative

Agricultural & Dairy Society Limite

Good safety on white clover, there is an initial check on growth, but it is quickly outgrown

•	Stock grazing can resume 7 days after application, there is a 21-day cutting interval Rainfast in 1 hour Apply at the rates and timings specified in the table below Each ProClova XL co-pack treats 3 hectares (7.4 acres) of newly sown leys or 2 hectares (5 acres) when applying	Сгор	Maximum Individual Dose	Application Timing
•		Newly Sown Leys (grass is less than 1 year old or less than 90% ground cover)	85 g/ha ProClova & 0.165 L/ha XL	From 3 true leaves of the grass, once the clover has reached the 3-trifoliate leaf stage and when weeds are actively growing. Apply from 1st April to 31st October.
•	to established grassland This product can also be applied at a higher use of rate once the newly sown ley has reached 90% ground cover	Established grassland (more than 1 year old or greater than 90% ground cover)	125 g/ha ProClova & 0.25 L/ha XL	When weeds are actively growing. Apply from 1st March to 31st October.







